

Hup Antibody References

Contents

Hup Antibody Isolation and Characterization.....	2
Studies Utilizing Hup Antibodies.....	2
Hup 1N	2
Hup 2B.....	4
Hup 1M	5
Hup1N and Hup2B	5
Hup 1M and Hup2B.....	7

Updated April 2, 2024

Hup Antibody Isolation and Characterization

1. Stanker LH, Wyrobek A, and Balhorn R; Monoclonal antibodies to human protamines. *Hybridoma* **3**: 293-303 (1987).
2. Stanker LH, McKeown C, Balhorn R, Lee C, Mazrimas J, Goralka M, and Wyrobek A; Immunological evidence for a P2 protamine precursor in mature rat sperm. *Mol Reprod Devel* **33**: 481-488 (1992).
3. Stanker LH, Wyrobek A, McKeown C, and Balhorn R; Identification of the binding site of two monoclonal antibodies to human protamine. *Mol Immunol* **18**: 1633-8 (1993).

Studies Utilizing Hup Antibodies

Hup 1N

1. Giorgini F, Davies HG, and Braun RE; Translational repression by MSY4 inhibits spermatid differentiation in mice. *Development* **129**: 3669-3679 (2002).
2. Sarmento OF, Digilio LC, Wang Y, Perlin J, Herr JC, Allis CD and Coonrod SA; Dynamic alterations of specific histone modifications during early murine development. *J Cell Sci* **117**: 4449-4459 (2004).
3. Aleem M, Choudhari J, Padwal V, Balasinor N, Parte P and Gill-Sharma MK; Hyperprolactinemia affects spermiogenesis in adult male rats. *J Endocrinol Investig* **28**(1): 39-48 (2005).
4. Aleem M, Padwal V, Choudhari J, Balasinor N, Parte P and Gill-Sharma MK; Tamoxifen affects expression of transition proteins and protamine genes in adult rats adversely influencing the sperm DNA condensation. *Asian J Androl* **7**(3): 311-321 (2005).
5. Aleem M, Padwal V, Choudhari J, Balasinor N, Parte P and Gill-Sharma MK; Cyproterone acetate affects protamine gene expression in the testis of adult male rat. *Contraception* **71**(5): 379-391 (2005).
6. Aleem M, Padwal P, Choudhari J, Balasinor N, Parte P and Gill-Sharma MK; Estradiol affects Androgen-Binding Protein expression in the testis of adult male rats. *Mol Cell Endocrinol* **253**: 1-13 (2006).
7. Aoki VW, Emery BR, Liu L and Carrell DT. Protamine levels vary between individual sperm cells of infertile human males and correlate with viability and DNA integrity. *J Androl* **27**: 890 (2006)
8. Zhang X, Gabriel MS and Zini A; Sperm nuclear histone to protamine ratio in fertile and infertile men: Evidence of heterogeneous subpopulations of spermatozoa in the ejaculate. *J Androl* **27**: 414 (2006).
9. Aleem M, Padwal V, Choudhary J, Balasinor N and Gill-Sharma MK; Sperm protamine levels as indicators of fertilizing potential in sexually mature rats. *Andrologia* **39**: 1-9 (2007).
10. Gill-Sharma MK; Prolactin and male fertility: The long and short feedback regulation (Review Article). *Int J Endocrinol Vol 2009* (article ID 687259):1-13 (2008).
11. Codrington AM, Hales BF and Robaire B; Exposure of male rats to cyclophosphamide alters the chromatin structure and basic proteome in spermatozoa. *Hum Reprod* **22**: 1431-42 (2007)
12. Ramos L, van der Heijden GW, Derijk A, Berden JH, Kremer JAM, van der Vlag J and de Boer P; Incomplete nuclear transformation of human spermatozoa in oligo-astheno-teratospermia: characterization by indirect immunofluorescence of chromatin and thiol status. *Hum Reprod* **23**: 259-270 (2008).
13. Worawittayawong P, Leigh C, Weerachatyanukul W, Manochantr S, Sobhon R, Breed WG and Sretarugsa P; Changes in distribution of basic nuclear proteins and chromatin organization

- during Spermiogenesis in the greater bandicoot rat, *Bandicota indica*. *Cell Tissue Res* **334**: 135-144 (2008).
14. Suphamungmee W, Wanichanon C, Vanichvirizakit R and Sobhon P; Spermiogenesis and chromatin condensation in the common tree shrew, *Tupaia glis*. *Cell Tissue Res*, **331**: 687-699 (2008).
 15. Yanagiva A, Delbes G, Svitkin YV, Robaire B and Sonenberg N; The poly(A)-binding protein Paip2a controls translation during late Spermiogenesis in mice. *J Clin Invest*, **120**(9): 3389-3400 (2010).
 16. Abou-Elhamd, A. S.; Sumayli, S.; Steger, K.; Ali, A. K. M.; Zayed, A. E. Effect of Khat (*Catha edulis* Forsk) extract on testicular maturation in pre-pubertal and pubertal rats: A morphological and biochemical study. *Anat Histol Embryol* 2021, 50 (2), 271-283. DOI: 10.1111/ahe.12626 From NLM.
 17. Codrington, A. M.; Hales, B. F.; Robaire, B. Exposure of male rats to cyclophosphamide alters the chromatin structure and basic proteome in spermatozoa. *Hum Reprod* 2007, 22 (5), 1431-1442. DOI: dem002 [pii] 10.1093/humrep/dem002.
 18. Gill, M. E.; Kohler, H.; Peters, A. Dual DNA staining enables isolation of multiple sub-types of post-replicative mouse male germ cells. *Cytometry A* 2022, 101 (6), 529-536. DOI: 10.1002/cyto.a.24539 From NLM.
 19. Gill-Sharma, M. K.; Choudhuri, J.; D'Souza, S. Sperm chromatin protamination: an endocrine perspective. *Protein Pept Lett* 2011, 18 (8), 786-801. DOI: BSP/ PPL/ E pub/0320 [pii].
 20. Gou, L. T.; Lim, D. H.; Ma, W.; Aubol, B. E.; Hao, Y.; Wang, X.; Zhao, J.; Liang, Z.; Shao, C.; Zhang, X.; et al. Initiation of Parental Genome Reprogramming in Fertilized Oocyte by Splicing Kinase SRPK1-Catalyzed Protamine Phosphorylation. *Cell* 2020, 180 (6), 1212-1227.e1214. DOI: 10.1016/j.cell.2020.02.020 From NLM.
 21. Hammadeh, M. E.; Hamad, M. F.; Montenarh, M.; Fischer-Hammadeh, C. Protamine contents and P1/P2 ratio in human spermatozoa from smokers and non-smokers. *Hum Reprod* 2010, 25 (11), 2708-2720. DOI: 10.1093/humrep/deq226 [pii].
 22. Hammoud, S.; Liu, L.; Carrell, D. T. Protamine ratio and the level of histone retention in sperm selected from a density gradient preparation. *Andrologia* 2009, 41 (2), 88-94. DOI: 10.1111/j.1439-0272.2008.00890.x
 23. Kobayashi, Y.; Tomizawa, S. I.; Ono, M.; Kuroha, K.; Minamizawa, K.; Natsume, K.; Dizdarević, S.; Dočkal, I.; Tanaka, H.; Kawagoe, T.; et al. Tsga8 is required for spermatid morphogenesis and male fertility in mice. *Development* 2021, 148 (8). DOI: 10.1242/dev.196212 From NLM.
 24. Miura, K.; Matoba, S.; Hirose, M.; Ogura, A. Generation of chimeric mice with spermatozoa fully derived from embryonic stem cells using a triple-target CRISPR method for Nanos3^t. *Biol Reprod* 2021, 104 (1), 223-233. DOI: 10.1093/biolre/ioaa176 From NLM.
 25. Noblanc, A.; Damon-Soubeyrand, C.; Karrich, B.; Henry-Berger, J.; Cadet, R.; Saez, F.; Guiton, R.; Janny, L.; Pons-Rejraji, H.; Alvarez, J. G.; et al. DNA oxidative damage in mammalian spermatozoa: where and why is the male nucleus affected? *Free Radic Biol Med* 2013, 65, 719-723. DOI: 10.1016/j.freeradbiomed.2013.07.044
 26. Ren, S.; Yang, D.; Dong, Y.; Ni, W.; Wang, M.; Xing, L.; Liu, T.; Hou, W.; Sun, W.; Zhang, H.; et al. Protamine 1 as a secreted colorectal cancer-specific antigen facilitating G1/S phase transition under nutrient stress conditions. *Cell Oncol (Dordr)* 2023, 46 (2), 357-373. DOI: 10.1007/s13402-022-00754-w From NLM.
 27. Suphamungmee, W.; Wanichanon, C.; Vanichviriyakit, R.; Sobhon, P. Spermiogenesis and chromatin condensation in the common tree shrew, *Tupaia glis*. *Cell Tissue Res* 2008, 331 (3), 687-699. DOI: 10.1007/s00441-007-0557-5 From NLM.

28. Yanagiya, A.; Delbes, G.; Svitkin, Y. V.; Robaire, B.; Sonenberg, N. The poly(A)-binding protein partner Paip2a controls translation during late spermiogenesis in mice. *J Clin Invest* 2010, **120** (9), 3389-3400. DOI: 10.1172/JCI43350.
29. Yuen, B. T.; Bush, K. M.; Barrilleaux, B. L.; Cotterman, R.; Knoepfler, P. S. Histone H3.3 regulates dynamic chromatin states during spermatogenesis. *Development* 2014, **141** (18), 3483-3494. DOI: 10.1242/dev.106450.
30. Zhou, J.; Yang, F.; Leu, N. A.; Wang, P. J. MNS1 is essential for spermiogenesis and motile ciliary functions in mice. *PLoS Genet* 2012, **8** (3), e1002516. DOI: 10.1371/journal.pgen.1002516 PGENETICS-D-11-01820 [pii].
31. Zhou, Q.; Xu, M.; Wang, X.; Yu, M.; Chen, X.; Lu, J.; Zhou, R.; Zhang, J.; Ling, X.; Ji, J. Deficiency of TBL1XR1 causes asthenozoospermia. *Andrologia* 2021, **53** (3), e13980. DOI: 10.1111/and.13980 From NLM.

Hup 2B

1. Bunick D, Balhorn R, Stanker LH and Hecht NB; Expression of the rat Protamine 2 gene is suppressed at the level of transcription and translation. *Exp Cell Res* **188**: 147-152 (1990).
2. Unni E, Zhang Y, Meistrich ML and Balhorn R; Rat spermatid basic nuclear protein TP3 is the precursor of Protamine 2. *Exp Cell Res* **210**: 39-45 (1994).
3. Lee K, Haugen HS, Clegg CH and Braun RE; Premature translation of protamine 1 mRNA causes precocious nuclear condensation and arrests spermatid differentiation in mice. *Proc Natl Acad Sci USA* **92**: 12451-12455 (1995).
4. Bench G, Corzett MH, DeYebra L, Oliva R and Balhorn R; Protein and DNA contents in sperm from an infertile human male possessing protamine defects that vary over time. *Mol Reprod Devol* **50**: 345-53 (1998).
5. Giorgini F, Davies HG and Braun RE; Translational repression by MSY4 inhibits spermatid differentiation in mice. *Development* **129**: 3669-3679 (2002).
6. Zhang X, Gabriel MS and Zini, A; Sperm nuclear histone to protamine ratio in fertile and infertile men: Evidence of heterogeneous subpopulations of spermatozoa in the ejaculate. *J Androl* **27**: 414 (2006).
7. Aoki VW, Emery BR, Liu L and Carrell DT; Protamine levels vary between individual sperm cells of infertile human males and correlate with viability and DNA integrity. *J Androl* **27**: 890 (2006).
8. Ramos L, van der Heijden GW Derijk A, Berden JH, Kremer JAM, van der Vlag J and de Boer P. Incomplete nuclear transformation of human spermatozoa in oligo-astheno-teratospermia: characterization by indirect immunofluorescence of chromatin and thiol status. *Hum Reprod* **23**: 259-270 (2008).
9. Yamauchi Y, Riel JM, Stoytcheva Z, Burgoyne PS and Ward MA, Deficiency in mouse Y chromosome long arm gene complement is associated with sperm DNA damage. *Genome Biology* **11**:R66 (2010).
10. de Mateo, S.; Ramos, L.; de Boer, P.; Meistrich, M.; Oliva, R. Protamine 2 precursors and processing. *Protein Pept Lett* 2011, **18** (8), 778-785. DOI: BSP/ PPL/ E pub/0319.
11. Eckhardt, M.; Wang-Eckhardt, L. A commercial human protamine-2 antibody used in several studies to detect mouse protamine-2 recognizes mouse transition protein-2 but not protamine-2. *Mol Hum Reprod* 2015, **21** (11), 825-831. DOI: 10.1093/molehr/gav046gav046.
12. Schneider, S.; Balbach, M.; Jan, F. J.; Fietz, D.; Nettersheim, D.; Jostes, S.; Schmidt, R.; Kressin, M.; Bergmann, M.; Wachten, D.; et al. Re-visiting the Protamine-2 locus: deletion, but not haploinsufficiency, renders male mice infertile. *Sci Rep* 2016, **6**, 36764. DOI: 10.1038/srep36764.

Hup 1M

13. Wyrobek AJ, Currie M, Stilwell JL, Balhorn R and Stanker LH; Detecting specific-locus mutations in human sperm. In: *Banbury Report 34: Biology of Mammalian Germ Cell Mutagenesis*, Cold Spring Harbor Laboratory Press, pp 93-111 (1990)
14. Bench G, Corzett MH, DeYebra L, Oliva R and Balhorn R; Protein and DNA contents in sperm from an infertile human male possessing protamine defects that vary over time. *Mol Reprod Devel* **50**: 345-53 (1998).
15. Li, Y, Lalancette, C, Miller, D and Krawetz, SA. Characterization of nucleohistone and nucleoprotamine components in the mature human sperm nucleus. *Asian J Androl* **10**: 535-541 (2008).

Hup1N and Hup2B

16. Zhong J, Peters AHFM, Keesock L and Braun RE; A double-stranded RNA binding protein required for activation of repressed messages in mammalian germ cells. *Nature Genetics*, **22**: 171-174 (1999).
17. Cho C, Willis WD, Goulding EH, Hung-Ha, H, Choi, Y-C, Hecht NB, and Eddy EM; Haploinsufficiency of protamine-1 or -2 causes infertility in mice. *Nature Genetics* **28**: 82-86 (2001).
18. Kimura T, Ito C, Watanabe S, Takahashi T, Ikawa M, Yomogida K, Fujita Y, Ikeuchi M, Asada N, Matsumiya K, Okuyama A, Okabe M, Toshimori K, and Nakano T; Mouse germ cell-less as an essential component for nuclear integrity. *Molecular and Cellular Biology*, **23**(4): 1304-1315 (2003).
19. Yan W, Ma L, Burns KH and Matzuk MM; HILS1 is a spermatid-specific linker histone H1-like protein implicated in chromatin remodeling during mammalian Spermiogenesis. *PNAS*, **100**(18): 10546-51 (2003).
20. Zhao M, Shirley CR, Mounsey S and Meistrich ML; Nucleoprotein transitions during spermiogenesis in mice and with transition nuclear protein Tnp1 and Tnp2 mutations. *Biology of Reproduction* **71**: 1016-1025 (2004).
21. Zhao M, Shirley CR, Hayashi S, Marcon L, Mohapatra B, Suanuma R, Behringer RR, Boissonneault G, Yanagimachi R and Meistrich ML; Transition nuclear proteins are required for normal chromatin condensation and functional sperm development. *Genesis*, **38**: 200-213 (2004).
22. Tanaka H, Iguchi N, Isotani A, Kitamura K, Toyama Y, Matsuoka Y, Onishi M, Masai K, Maekawa M, Toshimori K, Okabe M and Nishimune Y; HANP1/S1T2, a novel histone H1-like protein involved in nuclear formation and sperm fertility. *Molecular and Cellular Biology*, **25**(16): 7107-7119 (2005).
23. Hammoud S, Liu L and Carrell DT; Protamine ratio and the level of histone retention in sperm selected from a density gradient preparation. *Andrologia*, **41**(2): 88-94 (2009).
24. Hammadeh ME, Hamad MF, Montenarh M, Fischer-Hammadeh C; Protamine contents and P1/P2 ratio in human spermatozoa from smokers and non-smokers. *Hum. Reprod.* **25**(11):2708-2720 doi: 10.1093/humrep/deq226 (2010).
25. Abe, H.; Meduri, R.; Li, Z.; Andreassen, P. R.; Namekawa, S. H. RNF8 is not required for histone-to-protamine exchange in spermiogenesis. *Biology of Reproduction* 2021, 105 (5), 1154-1159. DOI: 10.1093/biolre/ioab132 (acccesed 4/2/2024).
26. Barral, S.; Morozumi, Y.; Tanaka, H.; Montellier, E.; Govin, J.; de Dieuleveult, M.; Charbonnier, G.; Coute, Y.; Puthier, D.; Buchou, T.; et al. Histone Variant H2A.L.2 Guides Transition Protein-Dependent Protamine Assembly in Male Germ Cells. *Mol Cell* 2017, 66 (1), 89-101 e108. DOI: S1097-2765(17)30162-4 [pii] 10.1016/j.molcel.2017.02.025.

27. Sophie Barral. Ph.D. Thesis, L'organisation post-méiotique de l'épigénome mâle. Biochimie, Biologie Moléculaire. Université Grenoble Alpes, May 25, 2016, 196 pages. Français. NNT : 2018GREAV058 <https://theses.hal.science/tel-02103249>.
28. Barton-Maclaren, T. S. Ph.D. Thesis, THE IMPACT OF CYCLOPHOSPHAMIDE ON MALE GERM CELL QUALITY AND CONSEQUENCES ON EARLY POST-FERTILIZATION EVENTS. McGill University, Montreal, Quebec, Canada, August 2007, 266 pages. Library and Archives Canada, Published Heritage Branch, ISBN: 978-0-494-50774-2.
29. de Vries, M.; Ramos, L.; Lacroix, R.; D'Hauwers, K.; Hendriks, J. C.; Kremer, J. A.; van der Vlag, J.; de Boer, P. Chromatin remodelling initiation in spermatids: differences among human males. *Andrology* 2013, 1 (3), 421-430. DOI: 10.1111/j.2047-2927.2013.00079.x.
30. Derijck, A. Ph.D. Thesis, The transmission of chromatin and DNA lesions by sperm and their fate in the zygote. Radboud University Nijmegen, the Netherlands, 2007, 350 pages, ISBN: 978-90-9022243-1, <http://hdl.handle.net/2066/52741>.
31. Dolan, C. E. Ph.D. thesis, Structural and Functional Studies of the Protamine 2-Zinc Complex from Syrian Gold Hamster (*Mesocricetus auratus*) Spermatids and Sperm. University of California, Davis, United States, September 16, 2004, 192 pages, UCRL-TH-206583.
32. El Zowalaty, A. E.; Baumann, C.; Li, R.; Chen, W.; De La Fuente, R.; Ye, X. Seipin deficiency increases chromocenter fragmentation and disrupts acrosome formation leading to male infertility. *Cell Death Dis* 2015, 6, e1817. DOI: 10.1038/cddis.2015.188 [pii].
33. El Zowalaty, A. E., Ph.D. Thesis, ESSENTIAL ROLES AND MECHANISMS OF SEIPIN IN MOUSE REPRODUCTION. University of Georgia, United States, Athens. Georgia, August 2017, 120 pages.
34. Giorgini, F.; Davies, H. G.; Braun, R. E. Translational repression by MSY4 inhibits spermatid differentiation in mice. *Development* 2002, 129 (15), 3669-3679.
35. Gunther, K.; Paradowska-Dogan, A.; Barmann, B.; Klein, H.; von Eichel-Streiber, C.; Hartley, R.; Weidner, W.; Behr, R.; Steger, K. Expression of sperm-specific protamines impairs bacterial and eukaryotic cell proliferation. *Histochem Cell Biol* 2015, 143 (6), 599-609. DOI: 10.1007/s00418-015-1309-3.
36. Jha, K. N.; Tripurani, S. K.; Johnson, G. R. TSSK6 is required for gammaH2AX formation and the histone-to-protamine transition during spermiogenesis. *J Cell Sci* 2017, 130 (10), 1835-1844. DOI: 10.1242/jcs.202721.
37. Kimura, T.; Ito, C.; Watanabe, S.; Takahashi, T.; Ikawa, M.; Yomogida, K.; Fujita, Y.; Ikeuchi, M.; Asada, N.; Matsumiya, K.; et al. Mouse germ cell-less as an essential component for nuclear integrity. *Mol Cell Biol* 2003, 23 (4), 1304-1315.
38. Li, W.; Wu, J.; Kim, S. Y.; Zhao, M.; Hearn, S. A.; Zhang, M. Q.; Meistrich, M. L.; Mills, A. A. Chd5 orchestrates chromatin remodelling during sperm development. *Nat Commun* 2014, 5, 3812. DOI: 10.1038/ncomms4812.
39. Li, W. The chromatin remodeler Chd5 orchestrates repackaging of the sperm genome Stony Brook University, December 2013, 146 pages.
40. Luense, L. J.; Donahue, G.; Lin-Shiao, E.; Rangel, R.; Weller, A. H.; Bartolomei, M. S.; Berger, S. L. Gcn5-Mediated Histone Acetylation Governs Nucleosome Dynamics in Spermiogenesis. *Dev Cell* 2019, 51 (6), 745-758.e746. DOI: 10.1016/j.devcel.2019.10.024 From NLM.
41. Merges, G. E.; Meier, J.; Schneider, S.; Kruse, A.; Fröbius, A. C.; Kirfel, G.; Steger, K.; Arévalo, L.; Schorle, H. Loss of Prm1 leads to defective chromatin protamination, impaired PRM2 processing, reduced sperm motility and subfertility in male mice. *Development* 2022, 149 (12). DOI: 10.1242/dev.200330 From NLM.
42. Ramos, L.; van der Heijden, G. W.; Derijck, A.; Berden, J. H.; Kremer, J. A.; van der Vlag, J.; de Boer, P. Incomplete nuclear transformation of human spermatozoa in oligo-astheno-

- teratospermia: characterization by indirect immunofluorescence of chromatin and thiol status. *Hum Reprod* 2008, 23 (2), 259-270. DOI: dem365 [pii] 10.1093/humrep/dem365.
43. Shiota, H.; Barral, S.; Buchou, T.; Tan, M.; Couté, Y.; Charbonnier, G.; Reynoird, N.; Boussouar, F.; Gérard, M.; Zhu, M.; et al. Nut Directs p300-Dependent, Genome-Wide H4 Hyperacetylation in Male Germ Cells. *Cell Rep* 2018, 24 (13), 3477-3487.e3476. DOI: 10.1016/j.celrep.2018.08.069 From NLM.
 44. Torres-Flores, U.; Díaz-Espinosa, F.; López-Santaella, T.; Rebollar-Vega, R.; Vázquez-Jiménez, A.; Taylor, I. J.; Ortiz-Hernández, R.; Echeverría, O. M.; Vázquez-Nin, G. H.; Gutierrez-Ruiz, M. C.; et al. Spermiogenesis alterations in the absence of CTCF revealed by single cell RNA sequencing. *Front Cell Dev Biol* 2023, 11, 1119514. DOI: 10.3389/fcell.2023.1119514 From NLM.
 45. van der Heijden, G. W.; Derijck, A. A.; Ramos, L.; Giele, M.; van der Vlag, J.; de Boer, P. Transmission of modified nucleosomes from the mouse male germline to the zygote and subsequent remodeling of paternal chromatin. *Dev Biol* 2006, 298 (2), 458-469. DOI: S0012-1606(06)00976-6 [pii] 10.1016/j.ydbio.2006.06.051.
 46. Yamaguchi, K.; Hada, M.; Fukuda, Y.; Inoue, E.; Makino, Y.; Katou, Y.; Shirahige, K.; Okada, Y. Re-evaluating the Localization of Sperm-Retained Histones Revealed the Modification-Dependent Accumulation in Specific Genome Regions. *Cell Rep* 2018, 23 (13), 3920-3932. DOI: 10.1016/j.celrep.2018.05.094 From NLM.
 47. Yelumalai, S., Ph.D thesis, Potential effects of assisted reproductive technology upon the abundance and localisation of two vital sperm proteins. University of Oxford, Jesus College, 2015, 341 pages.
 48. Zhong, J.; Peters, A. H.; Lee, K.; Braun, R. E. A double-stranded RNA binding protein required for activation of repressed messages in mammalian germ cells. *Nat Genet* 1999, 22 (2), 171-174. DOI: 10.1038/9684.
 49. Zhou, Q.; Xu, M.; Wang, X.; Yu, M.; Chen, X.; Lu, J.; Zhou, R.; Zhang, J.; Ling, X.; Ji, J. Deficiency of TBL1XR1 causes asthenozoospermia. *Andrologia* 2021, 53 (3), e13980. DOI: 10.1111/and.13980 From NLM.

Hup 1M and Hup2B

50. deYebra L, Ballesca J-L, Vanrell JA, Corzett M, Balhorn R and Oliva R; Detection of P2 precursors in the sperm cells of infertile patients who have reduced P2 levels. *Fertility and Sterility*, **69**(4): 755-9 (1998).
51. Carrell DT and Liu L; Altered protamine 2 expression is uncommon in donors of known fertility, but common among men with poor fertilizing capacity, and may reflect other abnormalities of spermiogenesis. *J Androl* **22**: 604-610 (2001).
52. Katayose H, Yanagida K, Hashimoto S, Yamada H, and Sato A; Use of diamide-acridine orange fluorescence staining to detect protamination of human-ejaculated sperm nuclei. *Fertil Steril*. **79**: Suppl 1: 670-6 (2003).
53. Katayose H, Takayama T, Hayashi S and Sato A; Role of mammalian sperm nuclear structure in fertilization and embryo development (Review Article). *Reproductive Medicine and Biology*, **5**(3): 161-168 (2006).